

NOTES ON NEW AND NOTEWORTHY PLANTS. CXXIII

Harold N. Moldenke

**LIPPIA BOLIVIANA** var. **INTEGRIFOLIA** Mold., var. nov.

Haec varietas a forma typica speciei foliorum laminis marginaliter integris recedit.

This variety differs from the typical form of the species in having the margins of its leaf-blades entire.

The type of the variety was collected by Adolfo M. Jiménez (no. 188) at Jatumpata, 25 km. from the city of Cochabamba, Prov. Tarata, Cochabamba, Bolivia, at 2590 meters altitude, on November 18, 1978, and is deposited in my personal herbarium. The collector comments: "Planta lenosa, 1—1.5 m. de altura, flor blanca".

-----

**ADDITIONAL NOTES ON THE ERIOCAULACEAE. LXXXIV**

Harold N. Moldenke

**SYNGONANTHUS GRACILIS** var. **RECURVIFOLIUS** Ruhl.

Additional bibliography: Moldenke, *Phytologia* 42: 44. 1979.

Santos describes this plant as 15 cm. tall, with "white" flowers, and found it growing on "original campo", flowering in August.

Additional citations: BRAZIL: Bahia: Santos 2365 (Z).

**SYNGONANTHUS GRACILIS** var. **TENUISSIMUS** Ruhl.

Additional bibliography: Moldenke, *Phytologia* 37: 76—77. 1977.

Hatschbach has found this plant growing in wet sandy soil near roadsides, flowering and fruiting in April.

Additional citations: BRAZIL: Minas Gerais: Hatschbach 41254 (Z).

**SYNGONANTHUS HATSCHBACHII** Moldenke

Additional bibliography: Moldenke, *Phytologia* 37: 79. 1977.

This species is obviously very closely related to S. harleyi Moldenke and may eventually prove to be only a form or variety of it.

Recent collectors describe the species as "crescendo em espaços abertos, folhas com linhas brancas nas margens" and have found it growing in restinga, flowering and fruiting in September.

Additional citations: BRAZIL: Bahia: Hatschbach 39668 (N—iso-type); Mori, Mattos Silva, & Santos 10471 (Ld). MOUNTED ILLUSTRATIONS: Moldenke, *Phytologia* 39: 237 & 238. 1978 (Z).